



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 26, 2015

Jonathan A. Janis
Regulatory Manager, NAFTA
Isagro USA, Inc
430 Davis Drive, Suite 240
Morrisville, NC 27560

Subject: Label Amendment – Numerous minor edits to the Master Label
Product Name: Badge X2
EPA Registration Number: 80289-12
Application Date: 4/6/2015
Decision Number: 503704

Dear Mr. Janis:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Please note that we did not review or approve the submitted Supplemental labels. In harmony with our routine procedure, Supplemental labels are processed mainly for new uses of the active ingredient, not for uses that have been on the label or me-too labels for three or more years.

If you have any questions, please contact Lindsay Roe by phone at 703-347-0506, or via email at roe.lindsay@epa.gov.

Sincerely,


A handwritten signature in black ink, appearing to read "Tony Kish". The signature is fluid and cursive, with the first name "Tony" and last name "Kish" clearly distinguishable.

Tony Kish, Product Manager 22
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Badge® X₂

Fungicide/Bactericide for Agricultural Use



ACTIVE INGREDIENTS:

Copper Oxychloride (CAS No. 1332-40-7)*	23.82%
Copper Hydroxide (CAS No. 20427-59-2)*	21.49%

OTHER INGREDIENTS: 54.69%

TOTAL: 100.00%

*Metallic Copper (Cu) Equivalent is 28.2% by weight

Badge X2 is a dry flowable fungicide/bactericide.

KEEP OUT OF REACH OF CHILDREN

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label find someone to explain it to you in detail.)

See inside booklet for additional Precautions and Directions For Use

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>NOTE TO PHYSICIAN: Possible mucosal damage may contraindicate use of gastric lavage. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.</p> <p>For Chemical Emergency Spill Leak Fire Exposure or Accident Call CHEMTREC Day or Night Domestic North America 800-424-9300 International 703-527-3887 (collect calls accepted)</p>	

EPA Registration No.: 80289-12

EPA Establishment No.:

Manufactured by:
 Isagro S.p.A. (d/b/a Isagro USA, Inc.)
 For:
 Isagro USA, Inc.
 430 Davis Drive, Suite 240
 Morrisville, NC 27560

 **FOR ORGANIC PRODUCTION**



Distributed by:
 Gowan Company
 P.O. Box 5569
 Yuma, AZ 85366-5569



Gowan
 The Go To Company

Made in Italy

NET CONTENTS:

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING/AVISO

May be fatal if swallowed. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if inhaled. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber, Selection Category A). Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators, and other handlers must wear the following:

- long-sleeved shirt and long pants
- shoes plus socks
- protective eyewear (goggles, safety glasses, or face shield)
- chemical-resistant gloves such as Natural Rubber

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff.

This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers, adults, children or pets, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Notify workers of the application by warning them orally.

GREENHOUSE USE; For at least 7 days following the application of copper-containing products in greenhouses:

- At least one container or station designed specifically for flushing eyes is available in operating

condition with the WPS-required decontamination supplies for workers entering the area treated with copper-containing products,

- Workers are informed orally, in a manner they can understand:
 - that residues in the treated area may be highly irritating to their eyes,
 - that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes,
 - that if they do get residues in their eyes, they should immediately flush their eyes with the eye flush container that is located with the decontamination supplies and
 - how to operate the eye flush container or eye flush station.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours for greenhouse uses and 48 hours for all other applications without required PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear if overhead exposure
- Protective eyewear (goggles, safety glasses, or face shield)

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter until sprays have dried.

INSTRUCTIONS

BADGE X₂ may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise in the specific crop use directions.

The per acre use rate of BADGE X₂ is applicable for both dilute and concentrate spraying. Depending upon the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to Minimum Recommended Spray Volume Table. Complete spray coverage is essential to ensure optimum performance from BADGE X₂. When treating by aerial application or with low volume application equipment, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

Consult the BADGE X₂ label for specific rates and timing of application by crop. **Where application rates and intervals are provided in a range (e.g. 4 to 12 pounds and 7 to 10 days), the higher rates and shorter spray intervals are recommended when rainfall is heavy and/or disease pressure is high. Use the higher rates for large mature tree crops.**

The pre harvest interval (PHI) for Badge X₂ is 0 days for all crops. Reentry into treated areas and harvest of treated crops can be performed when wearing required PPE during the 24 hour REI for greenhouse and 48 hour REI for field applications.

SPECIAL PRECAUTIONS

- BADGE X₂ **must not be applied** in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.
- Do not tank mix BADGE X₂ with Aliette® fungicide for use on any registered crops or ornamentals unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result. Use in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.
- This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, or other metallic surfaces.
- Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of BADGE X₂ resulting in possible phytotoxicity or loss of effectiveness.
- Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless directed on this label or by a state/local expert, it is advisable to test for compatibility and potential crop injury prior to commercial use of a new tank mix; otherwise do not tank mix.
- It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use.
- Do not apply this product through any irrigation (chemigation) system using aluminum parts or components as damage to the system may occur. Such application is prohibited regardless of whether the irrigation system is flushed with water after use of this product.
- Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s) which contain no aluminum parts or components. Do not apply this product through any other type of irrigation system.
- While volume is important in obtaining full spray coverage, often factors such as foliage density, environmental conditions and sprayer calibration have a greater impact. Always be sure that sprayers are calibrated to spray equipment manufacturer's specifications and environmental conditions are within those recommended by State and local regulatory authorities.
- When mixing, fill the spray tank one-half full with water. Add BADGE X₂ slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. **DO NOT PREMIX or SLURRY** BADGE X₂. Spreaders, stickers, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank or contact your chemical supplier. Observe all precautions and limitations on the labels of all products used in mixtures.

FROST INJURY PROTECTION (Bacterial Ice Nucleation Inhibitor)

Application of BADGE X₂ made to all crops listed on this label at the rates and stages of growth indicated, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola* and *Pseudomonas fluorescens*) and may therefore provide some protection against light frost. Not recommended for those geographical areas where weather conditions favor severe frost.

CROP USES

CITRUS: Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo and Tangerine.

FIELD CROPS: Alfalfa, Barley, Clover*, Corn, Oats, Peanut, Potato, Soybean, Sugar Beet, Sugarcane* and Wheat.

SMALL FRUITS: Blackberry, Blueberry, Cranberry, Currant, Gooseberry, Raspberry and Strawberry.

TREE CROPS: Almond, Apple, Apricot, Avocado, Banana, Cacao, Cherimoya, Cherry, Chestnut*, Coffee, Filbert, Mango, Nectarine, Nutmeg*, Olive, Peach, Pear, Pecan, Pistachio, Plantain, Plum, Prune, Quince and Walnut.

VEGETABLES: Artichoke, Asparagus*, Bean, Beet, Beet Greens, Broccoli, Brussels Sprout, Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac, Celery, Chard, Cucumber, Eggplant, Endive, Escarole, Garlic, Greens (Collard, Mustard and Turnip), Honeydew, Kale, Kohlrabi, Leek, Lettuce, Muskmelon, Okra, Onion, Pea, Pepper, Pumpkin, Radish, Rhubarb, Rutabaga, Shallot, Spinach, Squash, Tomato, Turnip, Watercress and Watermelon.

VINES: Grape, Hops and Kiwi.

MISCELLANEOUS: Atemoya, Carambola, Chives, Cilantro, Coriander, Dill, Ginseng, Guava, Litchi, Live Oak, Macadamia, Mamey Sapote, Mint, Papaya, Parsley, Passion Fruit, Persimmon, Rosemary, Sugar Apple and Sycamore, and Turfgrass*.

GREENHOUSE AND SHADEHOUSE CROPS: BADGE X₂ may be used in greenhouses and shadehouses to control diseases on any crop on this label where physiology allows greenhouse or shadehouse culture. While specific directions are presented for Citrus, Cucumber, Eggplant, Pepper, and Tomato; general use may occur for any crop on this label where physiology allows greenhouse or shadehouse culture.

ORNAMENTALS: Specified as listed.

*Not for use in California.

MINIMUM RECOMMENDED SPRAY VOLUME (GALLONS PER ACRE) WHEN APPLYING BADGE X ₂			
USE	AERIAL	DILUTE	CONCENTRATE*
Vegetables	3	20	—
Field Crops	3	20	—
Small Fruits	5	150	50
Vines	5	150	50
Tree Crops	10	400	50
Miscellaneous crops	10	150	50
Citrus	10	800	100
Ornamentals	10	100	50
*When using pesticide application equipment such as Curtec® or other similar sprayers which are capable of obtaining thorough coverage at low volumes, application rates as low as 20 gallons per acre of spray volume may be used.			

CROP USE DIRECTIONS

The following specific instructions are based on general application procedures. The recommendations of your local State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

CITRUS

BADGE X₂ may be mixed with dry foliar nutritionals (micronutrients) to create “Shot Bag” mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. BADGE X₂ per acre rates in these mixes must not exceed the maximum recommended label rates for disease control. Adding foliar nutritionals or other products to spray mixtures containing BADGE X₂ and applying to citrus during the post-bloom period when young fruit are present may result in spray burn.

DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MIN. RETREATMENT INTERVAL (DAYS)	COMMENTS
Algal Spot, Melanose, Scab	1.75 – 11**	45	12.6 ¹	7	Apply as pre-bloom and post-bloom sprays.
Greasy Spot, Pink Pitting	0.75 – 5**	45	12.6 ¹	7	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe.
Alternaria Brown Spot	1.75 – 7**	45	12.6 ¹	21	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to fruit should start after two thirds of the petals have fallen and be repeated on a 21 day schedule.
Phytophthora Brown Rot, Septoria Spot	1.75 – 7**	45	12.6 ¹	7	Begin application in fall before or just after the first rain and continue as needed. For brown rot only, apply to skirts of trees to a height of at least 4 feet. For control of septoria spot or where fruit have already been infected with brown rot, apply to entire tree. Apply also to bare ground 1 foot beyond skirt. NOTE: In California, in areas subject to copper injury, add 0.33 to 1 pound of high quality lime per pound of BADGE X ₂ .

DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MIN. RETREATMENT INTERVAL (DAYS)	COMMENTS
Phytophthora Foot Rot	0.5	45	12.6 ¹	7	Mix with 1 quart of water, Tre-Hold® or latex paint. Paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to 1 year, but does not cure existing infections. NOTE: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off.
Citrus Canker (Suppression)	1 – 11	45	12.6 ¹	7	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.
Black spot*	1 – 3**	45	12.6 ¹	7	Initiate treatment prior to or at the first appearance of disease and repeat on a 7 to 21 day interval as needed. Use short application intervals when conditions favor disease development.

NOTE: Phytotoxicity may occur on young tender flush when BADGE X₂ is applied to citrus seedlings grown in greenhouses or shadehouses.

¹Maximum annual amount allowed for all disease applications combined

* Not for use in California.

** Use the higher rates when conditions favor disease development.

CITRUS (FIELD NURSERY GROWN)

To control Melanose, Scab, Pink Pitting, Greasy Spot and Brown Rot and for suppression of Citrus Canker, apply 1.75 to 3.5 lbs of product per acre. Apply BADGE X₂ at 28 day intervals depending on disease severity and rainfall. The maximum single application rate is 3.15 lbs of Cu / A. The maximum annual application rate is 12.6 pounds of Cu / A. The minimum retreatment interval is 7 days.

FIELD CROPS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/Year (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	0.75 – 1.5	4.0	1.12	30	Apply 10 to 14 days before each harvest or earlier if disease threatens. NOTE: Spray injury may occur with sensitive varieties such as Lahontan.
Cereal Grains: (Barley, Millet, Oat, Rye, Sorghum, Wheat)	Fusarium Head Blight (Suppression), Helminthosporium, Powdery Mildew (Suppression), Septoria Leaf Blotch, Spot Blotch, Stagonospora Leaf and Glume Blotch, and Stem Rust	0.5 – 1.8**	3.8	1.06	10	BADGE X ₂ can be applied as a foliar application for early season disease control and again at early heading then followed with another application 10 days later.

FIELD CROPS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/Year (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Clover***	Anthracnose, Bacterial Blight, Bacterial Leaf Spot, Cercospora Leaf Spot, Powdery Mildew	0.5 – 1.8**	16.9	4.74	7	Begin applications when conditions first favor disease development and repeat at 7 to 14 day intervals.
Corn (Field Corn, Popcorn, Seed Corn, Sweet Corn)	Bacterial Stalk Rot	0.5* – 2.5**	15.0	4.2	7	Begin treatment when disease first appears and repeat every 7 to 10 days. Use shorter spray intervals when conditions favor disease development. * In California, use the low rate of 0.5 lbs/A only.
Peanut	Leaf Spot, Rust	1 – 2.5**	16.9	4.74	7	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 10 to 14 day intervals. Reduce sprays to 7 day intervals during humid weather.
Potato	Early Blight, Late Blight, Powdery Mildew	1 – 4**	89.3	25	5	Apply 1 to 2 pounds at 7 to 10 day intervals starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 4 lbs/A when disease is more severe. Under conditions of severe disease, control with BADGE X ₂ will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners.
Soybean	Bacterial Blight, Downy Mildew, Powdery Mildew	0.75 – 2.5**	16.9	4.74	7	For preventive applications, begin first application when plant height reaches 6 inches and repeat on a 7 to 14 day interval as needed.
Sugar Beet	Cercospora Leaf Spot, Powdery Mildew	1 – 4**	28.1	7.86	10	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals. Addition of a spreader/sticker is recommended. Do not tank mix with glyphosate. Potential for phytotoxicity exists if an application of glyphosate is made within 7 to 10 days of a Badge X ₂ application.

FIELD CROPS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/Year (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Sugarcane***	Rusts (brown and orange)	1.75	3.8	1.06	10	Recommended for tank mixture with other products registered for rust control. For suppression of rust, begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals. Addition of a spreader/sticker is recommended.
<p>* In California, use the low rate of 0.5 lbs/A only on corn.</p> <p>** Use the higher rates when conditions favor disease development.</p> <p>*** Not for use in California</p>						

SMALL FRUITS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Brambles - Blackberry Raspberry (Aurora, Boysen, Cascade, Chehalem, Logan, Marion, Santiam, Thornless Evergreen)	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	1.75 – 3.5	35.7	10 ¹	7	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. If needed, agricultural-type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	1 – 2.25	35.7	10 ¹	7	Apply when leaf buds begin to open and repeat when flower buds show white. If needed, agricultural-type spray oil may be added. NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue application if signs of crop injury appear.
Blueberry	Bacterial Canker*	1.5 – 7**	30.0	8.4 ¹	28	Make first application before rain falls and a second application 4 weeks later. * In California, use the high rate of 7 lbs/A only.
	Fruit Rot, Phomopsis Twig Blight	1 – 4**	30.0	8.4 ¹	7	Dormant Application: Begin applications when bloom buds begin to swell. Make additional applications at 10 to 14 day intervals before blooms open.
Cranberry	Fruit Rot	3.5 – 7	45.0	12.6 ¹	7	Make first application in late bloom. Apply one or two additional applications at 10 to 14 day intervals depending on disease severity.

SMALL FRUITS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
	Rose Bloom	3.5 – 7	45.0	12.6 ¹	7	Apply three sprays on 10 to 14 day schedule as soon as symptoms are observed.
	Bacterial Stem Canker	3.5 – 7	45.0	12.6 ¹	7	Apply postharvest and again in spring at bud swell. Apply one or two additional applications at 10 to 14 day intervals depending on disease severity.
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight (<i>Monilinia</i>)	3.5 – 7	45.0	12.6 ¹	7	Apply delayed dormant spray in the spring. Repeat at 10 to 14 day intervals through pre-bloom.
Currant, Gooseberry	Anthracnose, Leaf Spot	4.25 – 9	57.1	16	10	Make initial application after first leaves have expanded. Continue on a 10 to 14 day schedule during wet conditions in the spring. Make an additional application after harvest.
Strawberry	Angular Leaf Spot, (<i>Xanthomonas</i>), Leaf Blight, Leaf Scorch, Leaf Spot, Downy Mildew, Powdery Mildew	1 – 2.5**	29.2	8.19	7	Begin application when plants are established and continue on a weekly schedule throughout the season. Apply in at least 20 gallons of water. NOTE: Discontinue applications if signs of crop injury appear.
¹ Maximum annual amount allowed for all disease applications combined. * Blueberry- Bacteria Canker; In California, use the high rate of 7 lbs/A only. ** Use the higher rates when conditions favor disease development.						

TREE FRUIT						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Apple	Anthracnose, Blossom Blast, European Canker (<i>Nectria</i>), Shoot Blast (<i>Pseudomonas</i>)	5 – 14**	57.1	16 ¹	n/a	Apply before fall rains. Only 1 application per season is permitted. NOTE: Use on yellow varieties may cause discoloration. To avoid discoloration, pick before spraying.
	Apple Scab, Fire Blight (single application rate)	3.5 – 7	57.1	16 ¹	n/a	Make application between silver-tip and green-tip. Apply as a full-cover spray for early season disease suppression. Only 1 application per season is permitted. NOTE: Moderate to severe crop injury may occur from late application; discontinue use when green-tip reaches ½ inch.
	Apple Scab (extended spray application rate)	0.75 – 1.75	57.1	16 ¹	5	Extended spray schedule where fruit finish is not a concern: Continued

TREE FRUIT						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMEN T INTERVAL (DAYS)	COMMENTS
	Fire Blight (extended spray application rate)	0.5 – 1.5	57.1	16 ¹	5	application may be made at 5 to 7 day intervals between 0.5 inch green-tip and first cover spray. NOTE: Moderate to severe crop injury may result from this extended spray schedule. It is not intended for fresh market apples or fresh apples where fruit finish is a concern as it is likely to cause fruit russetting. The addition of 1 to 3 pounds of hydrated lime per pound of BADGE X ₂ may reduce crop injury.
	Bitter Rot, Black Spot, Blotch, Powdery mildew	1 – 2.8	57.1	16 ¹	5	Begin applications at petal fall and repeat through fourth cover spray. The addition of 3 to 5 lbs hydrated lime per 100 gallons may reduce crop injury.
	Brooks Spot	2	57.1	16 ¹	5	Apply Badge X ₂ plus 2 lbs hydrated lime per 100 gallons of water. Make applications during late cover sprays.
	Bullseye Rot	7.5	57.1	16 ¹	n/a	Use Badge X ₂ plus sprayable oil per 100 gallons water. Make application after harvest. Only one application per season.
	Collar Rot, Crown Rot	1 – 2.5	57.1	16 ¹	5	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply in early spring or in fall after harvest for best results. Do not apply to foliage or fruit. NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.
	Sooty Blotch	1 – 2.5	57.1	16 ¹	5	Use Badge X ₂ plus 2.5 lbs hydrated lime per 100 gallons water. Apply during late cover sprays. When conditions indicate the potential for increased copper injury, add additional lime.
Cherry, Plum, Prune	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Coryneum Blight (Shot Hole)	3.5 – 14**	64.3	18 ¹	7	Fall Applications: Make first application before fall rains and a second at late dormant (up to the pink bud stage). If needed, agricultural- type spray oil may be added. For cherries: Where disease is severe, an additional application shortly after harvest may be required.

TREE FRUIT						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	3.5 – 5**	64.3	18 ¹	5	Apply during early bloom. Do not apply after full bloom or injury may occur. Use the higher rates when rainfall is heavy.
	Black Knot (Plum)	1.5 – 5**	64.3	18 ¹	5	Make an application at bud swell up to early bloom for early season disease suppression. Apply before full bloom. Use the higher rates when rainfall is heavy and disease pressure is high. NOTE: To avoid plant injury, do not use after full bloom.
Cherry	Cherry Leaf Spot (Sour Cherries Only)	1 – 5	64.3	18 ¹	5	Cover Sprays: Apply at petal fall as well as 1 to 2 times after petal fall. Use the lower rates where disease infection is light and use the higher rates for a dormant application (up to the pink bud stage) or where disease infection is moderate to heavy. Do not apply to sweet cherry or the English Morello variety as severe injury will result. The addition of 1 to 3 pounds of hydrated lime per pound of BADGE X2 may reduce crop injury. NOTE: Moderate to severe injury such as leaf spotting and defoliation may occur from post-bloom applications.
Apricot, Nectarine, Peach	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Bacterial Spot, (<i>Xanthomonas</i>), Coryneum Blight (Shot Hole), Leaf Curl	3.5 – 14**	64.3	18 ¹	7	Fall Applications: Make first application before fall rains and a second at late dormant. For peach leaf curl, late dormant application must be made before leaf buds swell (up to the pink bud stage). Use the higher rates when rainfall is heavy. If needed, agricultural-type spray oil may be added.
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	3.5 – 5**	64.3	18 ¹	5	Full cover spray at pink bud.
	Bacterial Spot	0.5 – 2	64.3	18 ¹	5	Post-bloom application applied at first and second cover sprays. NOTE: Do not spray 3 weeks prior to harvest. Use only listed rates. Spotting of leaves and defoliation may occur from use in cover sprays.

TREE FRUIT						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Atemoya, Sugar Apple (Annona)	Anthrachnose	1.5 – 8**	45.0	12.6	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Avocado	Anthrachnose, Blotch, Scab	3.5 – 11**	67.5	18.9	14	Apply when bloom buds begin to swell and continue application at monthly intervals for 5 to 6 applications.
Banana, Plantain	Sigatoka (Black and Yellow)	0.75 - 2	67.5	18.9 ¹	7	Apply by air in 3 gallons of water. If needed, agricultural-type spray oil may be added. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.
	Black Pitting	1.75 – 3.5	67.5	18.9 ¹	7	Mix in 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
Carambola	Anthrachnose	2.5 – 7**	37.5	10.5	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Cherimoya (custard apple)	Anthrachnose	1 – 4**	30	8.4	14	Begin applications when conditions first favor disease development and repeat using a 14 day interval. Apply in sufficient water for thorough coverage. Make first application to a small area to test for crop sensitivity. The addition of 1 to 3 pounds of hydrated lime per 2 lbs of BADGE X ₂ may reduce crop injury.
Guava	Anthrachnose, Red Algae	1.25 – 4**	17.6	4.92	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Mamey Sapote	Algal Leaf Spot, Anthrachnose	2.5 – 7**	30.0	8.4	14	Apply when conditions favor disease development. Repeat on 14 to 30 day schedule as disease severity and environmental conditions dictate.
Mango	Anthrachnose	2 – 9**	171	48	7	Apply monthly after fruit set until harvest. Use the higher rates when rainfall is heavy.

TREE FRUIT						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Olive	Anthrachnose, Olive Knot, Olive Leaf Spot, Peacock Spot	3.5 – 11**	64.3	18	30	Make first application before winter rains begin. A second application in early spring should be made if disease is severe. Apply the higher rates for heavy disease pressure.
Papaya	Anthrachnose	2 – 9**	75.7	21.2	7	Apply before disease appears. Apply at 14 day intervals. The addition of an approved spreader is desirable.
Pear	Fire Blight	0.5 – 1	57.1	16 ¹	5	Apply at 5 day intervals throughout the bloom period. NOTE: Russetting may occur in copper sensitive varieties. Excessive dosages may cause fruit russet on any variety.
	Blossom Blast (<i>Pseudomonas</i>)	5.25 – 14**	57.1	16 ¹	n/a	Apply before fall rains and again during dormancy before spring growth starts. Use the higher rates when disease pressure is high. Only 1 application per season is permitted.
Persimmon	Cercospora Leaf Spot	0.75 – 1.5	21.4	6.0	14	Begin applications in May/June at leaf flush and repeat applications on a 14 day interval or greater depending on disease severity and environmental conditions.
Quince	Fire Blight, Blossom Blast	0.5 – 1	57.1	16	5	Apply at 5 day intervals throughout the bloom period. Apply in adequate water for thorough coverage.
¹ Maximum annual amount allowed for all disease applications combined						
** Use the higher rates when conditions favor disease development.						

TREE NUTS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Almond	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Coryneum Blight (Shot Hole)	3.5 – 14**	64.3	18 ¹	7	Fall Applications: Make first application before fall rains and a second at late dormant. If needed, agricultural-type spray oil may be added.
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	2.5 – 3.5**	64.3	18 ¹	5	Apply during early bloom. Do not apply after full bloom or injury may occur. Use the higher rates when rainfall is heavy.

TREE NUTS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
	Bacterial Blast (<i>Pseudomonas</i>)	0.5	64.3	18 ¹	5	Post-Bloom: To control bacterial blast in sprinkler irrigated orchards or when disease is severe, apply 0.5 lbs Badge X ₂ post-bloom at 2 week intervals or just prior to sprinkler irrigation. NOTE: Foliar injury may occur from post-bloom sprays on almonds, especially on NePlus varieties.
Cacao	Black Pod	1 – 7.5**	56.2	15.75	14	Begin applications at the start of the rainy season and continue while infection conditions persist. Apply 0.75 to 2 pounds at 14 to 21 day intervals depending on disease severity. For drier areas, make two to four applications using 2.5 to 3.75 lbs/A according to disease incidence and planting density.
Chestnut*	Leaf Spot	1 – 4	30	8.4	14	Begin applications when conditions first favor disease development. Make applications to protect shoot growth throughout the season. Use the lower rates where disease infection is light and use the higher rates for a dormant application or where disease infection is moderate to heavy.
Coffee	Coffee Berry Disease (<i>Colletotrichum coffeanum</i>)	2.5 – 7**	45.0	12.6 ¹	14	Apply first spray after flowering and before onset of long rains and then at 21 to 28 day intervals until picking.
	Bacterial Blight (<i>Pseudomonas syringae</i>)	2.5 – 7**	45.0	12.6 ¹	14	Begin spray program before the onset of long rainy periods and continue throughout the rainy season at 14 to 21 day intervals. The critical time of spraying to control this disease is just before, during and after flowering(s) especially when coinciding with wet weather. Use the higher rates when rainfall is heavy.
	Leaf Rust (<i>Hemileia vastatrix</i>)	1 – 3**	45.0	12.6 ¹	14	Apply before the onset of rain and then at 21 day intervals while the rains continue. Use the higher rates when rainfall is heavy.
	Iron Spot (<i>Cercospora coffeicola</i>), Pink Disease (<i>Corticium salmonicolor</i>)	0.75 – 1.5	45.0	12.6 ¹	28	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for 3 applications.

TREE NUTS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Filbert	Bacterial Blight	7 – 12**	85.7	24 ¹	14	Apply as a postharvest spray. In seasons of heavy rainfall apply a second spray when three-fourths of the leaves have dropped. Use the higher rates when rainfall is heavy. If needed, agricultural-type spray oil may be added. Use only in the states of Oregon and Washington.
	Eastern Filbert Blight	7 – 12**	85.7	24 ¹	14	Apply as a dilute spray in adequate water for thorough coverage. Make applications starting at bud swell to bud break and continue at 2 week intervals until early May. Thorough coverage is essential. Use the higher rates when rainfall is heavy. If needed, agricultural-type spray oil may be added. Use only in the states of Oregon and Washington.
Litchi	Anthracnose	1.25 – 4**	17.6	4.92	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
Macadamia	Anthracnose	2.5 – 8**	33.7	9.44 ¹	7	Initiate sprays at first sign of flowering and repeat on weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
	Phytophthora Blight (<i>P. capsici</i>), Raceme Blight (<i>Botrytis cinerea</i>)	1.25 – 5**	33.7	9.44 ¹	7	Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage.
Nutmeg*	Leaf Spot, Shot Hole	1 – 4	30	8.4	14	Begin applications when conditions first favor disease development. Make applications to protect leaves during the rainy season. Use the lower rates where disease infection is light and use the higher rates where disease infection is moderate to heavy. Make first application to a small area to test for crop sensitivity. The addition of 1 to 3 pounds of hydrated lime per 2 lbs of BADGE X ₂ may reduce crop injury.
Pecan	Kernel Rot, Shuck Rot (<i>Phytophthora cactorum</i>), Zonate Leaf Spot (<i>Cristulariella pyramidalis</i>)	1 – 4**	30.0	8.4 ¹	14	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to 4 week intervals starting at kernel growth and continue until shucks open. Use the shorter spray intervals if frequent rainfall occurs.

TREE NUTS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
	Ball Moss* Spanish Moss*	2.5 – 7	30.0	8.4 ¹	365	Apply in 100 gallons of water in the spring when ball moss is actively growing, using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. The addition of a nonionic surfactant will improve control. A second application may be required after 12 months.
Pistachio	Botryosphaeria Panicle and Shoot Blight, Botrytis Blight, Late Blight (<i>Alternaria alternata</i>), Septoria Leaf Blight	1.75 – 7**	30.0	8.4	14	Make initial application at bud swell and repeat on a 14 to 28 day schedule. If disease conditions are severe, use the shorter spray intervals.
Walnut	Walnut Blight	3.5 – 11**	114.3	32	7	Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage when frequent rainfall or extended periods of moisture occur. Thorough coverage of catkins, leaves and nutlets is essential for effective control. NOTE: Adequate control may not be obtained when copper tolerant species of Xanthomonas bacteria are present.
¹ Maximum annual amount allowed for all disease applications combined * Not for use in California ** Use the higher rates when conditions favor disease development.						

VEGETABLES						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/ YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Artichoke	Ramularia Leaf Spot, Powdery Mildew	0.5* – 1.75**	9.5	2.65	7	Recommended for tank mixture with other products registered for control of listed diseases. For suppression, begin applications when conditions first favor disease development and repeat using a 7 day interval. Addition of a spreader/sticker is recommended. * In California use the low rate of 0.5lbs/A only.

VEGETABLES						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/ YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Asparagus***	Rust	1 – 2.5**	17.6	5	10	Recommended for tank mixture with other products registered for control of rust. For suppression, begin applications when conditions first favor disease development and repeat using a 10 day interval. Addition of a spreader/sticker is recommended.
Bean (Dry, Green)	Anthrachnose, Bacterial Blight, Brown Spot, Common Blight, Cercospora Leaf Spot, Downy Mildew, Halo Blight	0.5 – 2**	16.9	4.74	7	For protective sprays, make first application when plants are 6 inches high; repeat on a 7 to 14 day schedule depending on environmental conditions.
Beet (Table Beet, Beet Greens)	Cercospora Leaf Spot, Downy Mildew	0.75 – 4**	28.1	7.86	10	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals.
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot, Downy Mildew	0.75 – 3.57**	17.9	5	7	Begin applications when disease first threatens and repeat at 7 to 14 day intervals depending on disease severity.
Celery, Celeriac	Bacterial Blight, Cercospora Early Blight, Downy Mildew, Septoria Late Blight	0.75 – 3.57	18.9	5.3	7	Begin applications as soon as plants are first established in the field, repeating at 7 day intervals depending on disease severity and environmental conditions.
Chard	Cercospora Leaf Spot, Ramularia Leaf Spot	0.5 – 2.5**	14.1	3.95	7	Begin applications when conditions first favor disease development and repeat at 7 to 14 day intervals.
Crucifers (broccoli, brussels sprout, cabbage, cauliflower, Chinese cabbage, collard greens, kale, kohlraabi mustard greens, turnip greens)	Black Leaf Spot (<i>Alternaria</i>), Black Rot (<i>Xanthomonas</i>), Downy Mildew	0.5 – 1.8**	9.5	2.65	7	Apply at 7 to 10 day intervals. Begin application after transplants are set in the field or shortly after emergence of field seeded crops or when conditions favor disease development. The addition of a spreader/sticker may enhance retention of spray deposition on cruciferous crops. NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.

VEGETABLES						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/ YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Cucurbits (cantaloupe, casaba, chayote, citron melon, cucumber, gourd honeydew, muskmelon, pumpkin, squash (summer and winter), watermelon and waxgourd, Chinese	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (Suppression)	0.5 – 2.5**	18.8	5.25	5	Begin applications prior to disease development and continue while conditions are favorable for disease development. Repeat sprays at 5 to 7 day intervals. NOTE: Crop injury may occur from application at higher rates and shorter intervals. Discontinue use if injury occurs.
Eggplant	Alternaria Blight, Anthracnose, Downy Mildew, Phomopsis, Phytophthora	0.75 – 1.5	28.2	7.9	7	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals depending on disease severity.
Lettuce (Head and Leaf), Endive, Escarole	Anthracnose, Downy Mildew, Leaf Spot, Powdery Mildew	0.75 – 1.5	28.6	8.0	5	Begin treatment at the first sign of disease. Repeat on a 7 to 10 day interval to suppress disease. Slight injury may occur under adverse conditions.
Okra	Anthracnose, Bacterial Leaf Spot, Leaf Spots, Pod Spot, Powdery Mildew	0.75 – 1.75**	18.8	5.25	5	Begin treatment when conditions are favorable for disease development and repeat using a 5 to 10 day interval as needed. Use shorter intervals when conditions favor disease.
Onion, Garlic, Leek, Shallot	Alternaria, Bacterial Blight, Downy Mildew, Purple Blotch, Rust	0.75 – 2.75	21.4	6	7	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals depending on disease severity. Can cause phytotoxicity to leaves.
Pea	Powdery Mildew	0.5 – 2.5**	14.1	3.95	7	Begin applications when disease symptoms first appear and repeat at weekly intervals.
Pepper (bell, chili)	Alternaria, Anthracnose, Bacterial Spot, Cercospora Leaf Spot, Downy Mildew, Early and Late Blight, Phytophthora Blight*	0.75 – 2.25**	42.3	11.85	3	Begin applications when conditions first favor disease development and repeat at 7 to 10 day intervals depending on disease severity.
Radish, Rutabaga, Turnip	Alternaria, Anthracnose, Bacterial Leaf Spot, Black Rot, Cercospora Leaf Spot, Downy Mildew, White Rust	1 – 2.25**	28.1	7.86	10	Begin application when disease first appears or when conditions favor disease development. Repeat using a 10 day interval.

VEGETABLES						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/ YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Rhubarb	Leaf Spot	1 – 2.25**	13.9	3.95	7	Begin application when disease first appears or when conditions favor disease development. Repeat at 7 to 10 day intervals.
Spinach	Anthracnose, Blue Mold, Cercospora Leaf Spot, Downy Mildew, White Rust	0.75 – 1.25**	14.1	3.95	7	Begin application when disease first appears or when conditions favor disease development. Repeat at 7 to 10 day intervals. NOTE: Flecking may occur on spinach leaves.
Tomato Fresh market	Anthracnose, Bacterial Canker, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	0.75 – 1.8**	26.6	6.0	3	Begin application when disease first threatens and repeat at 5 to 10 day intervals depending on disease severity.
Processing			62.1	17.4		
Watercress	Cercospora Leaf Spot	0.75 – 1.8	7.6	2.12	7	Begin applications when plants are first established in the field, repeating at 7 to 14 day intervals depending on disease severity. Do not exceed four applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.
* Artichoke- In California use the low rate of 0.5lbs/A only. ** Use the higher rates when conditions favor disease development. *** Not for use in California						

VINES						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Grape	Black Rot, Downy Mildew, Phomopsis, Powdery Mildew	0.75 – 3.5**	71.4	20	3	Begin applications at bud break with subsequent applications throughout the season depending on disease severity. NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara and Rosette. Either test for sensitivity or add 1 to 3 pounds of hydrated lime per pound of BADGE X ₂ .

VINES						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Hops	Downy Mildew	0.75 – 1.8	9.5	2.65	10	Make crown treatment after pruning, but before training. After training, additional treatments are needed at about 10 day intervals. NOTE: Discontinue use 2 weeks before harvest.
Kiwi	<i>Erwinia herbicola</i> , <i>Pseudomonas fluorescens</i> , <i>Pseudomonas syringae</i>	2 – 7	22.5	6.3	30	Apply in 200 gallons of water per acre. Make applications on a monthly basis. A maximum of three applications may be made.
Passion Fruit	Anthraxnose	2 – 8**	33.7	9.44	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage.
** Use the higher rates when conditions favor disease development.						

MISCELLANEOUS CROPS						
CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Chives	Downy Mildew	0.75 – 1.8	9.5	2.65	7	Begin applications when plants are established in the field. Repeat applications every 7 to 10 days depending on disease conditions.
Cilantro, Coriander, Rosemary	Leaf Spot	0.75* – 1.5	9.34	2.65	10	Begin applications when plants are established in the field. Begin applications prior to disease development and repeat every 10 days depending on disease conditions. * Rosemary- In California, use the low rate of 0.75 lbs/A only.
Dill	Phoma Leaf Spot, Rhizoctonia Foliage Blight	0.75 – 2.5**	14.1	3.95	7	Begin applications when plants are first established in the field and repeat at 7 to 10 day intervals depending upon disease severity and environmental conditions.

MISCELLANEOUS CROPS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Ginseng	Alternaria Leaf Blight, Stem Blight	1 – 3.5	18.8	5.25	7	Use as a tank mix with 2 pounds Rovral® 50W in 100 gallons of water. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Begin BADGE X ₂ -Rovral applications as soon as plants have emerged in spring. Application should be repeated every 7 days until plants become dormant in fall. Apply fungicides at least 8 hours before rain. Use of a spreader-sticker or sticker is advised. NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions such as those found in the dense canopies of 2 to 4 year old Ginseng. It is very important that the stems be thoroughly covered with fungicide; therefore, use a spray apparatus which distributes the fungicide throughout the canopy.
Live Oak	See ORNAMENTAL section for Control of Ball Moss*, Spanish Moss* and Lichens* on Ornamentals and Shade Trees.					
Mint	Downy Mildew, Leaf Spot, Powdery Mildew, Rust	0.75* – 1.5	9.34	2.65	10	Begin applications prior to disease development and repeat every 10 days depending on disease conditions. * In California, use the low rate of 0.75 lbs/A only.
Parsley	Bacterial Blight (<i>Pseudomonas</i> <i>sp.</i>)	1.25 – 2.8	7.1	2	10	Begin applications when plants are first established in the field and repeat at 10 day intervals as needed depending on disease severity and environmental conditions.
Sycamore	Anthrachnose	0.75 – 2.5**	12.9	3.6	7	Apply as a full cover spray in 100 gallons of water or sufficient volume for thorough coverage. Make first application at bud crack and second application 7 to 10 days later at 10% leaf expansion.

MISCELLANEOUS CROPS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	MAX. ANNUAL RATE (LBS Cu/A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Turfgrass***	Algae	4 – 6	75	21	10	May be used as a maintenance spray as needed. May be used in combination with other fungicides. Use a minimum application volume of 100 gallons of water per acre. Apply to a small area prior to large area applications to check for phytotoxicity. If phytotoxicity is present, discontinue use.

* Rosemary & Mint- In California, use the low rate of 0.75 lbs/A only.

** Use the higher rates when conditions favor disease development.

*** Not Registered for use in California.

GREENHOUSE AND SHADEHOUSE CROPS

Notice to User: BADGE X₂ may be used in greenhouses and shadehouses to control disease on crops which appear on this label and specific instructions have been developed for the crops listed. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shadehouses differs greatly from crops grown under field conditions. Neither the manufacturer nor seller has determined whether or not BADGE X₂ can be used safely on all greenhouse and shadehouse grown crops. The user should determine if BADGE X₂ can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., foliage, fruit, etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use.

Apply BADGE X₂ according to specific rates given for those crops in pounds per acre. Product contains 0.01 lbs of metallic copper per tablespoon. **One and a half (1.5) level tablespoons of BADGE X₂ per 1000 square feet is equivalent to 2.4 lbs/A.** BADGE X₂ should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat as labelled; use shorter spray intervals during periods when severe disease conditions persist.

CROP	DISEASE	APP. RATE (TBSP PRODUCT / 1000 SQ. FT.)	MAX. APP. RATE/YEAR (TBSP Product /1000 SQ. FT.)	MAX. ANNUAL RATE (pounds Cu/1000 SQ. FT.)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Citrus (Non-Bearing Nursery)	Brown Rot, Citrus Canker, Greasy Spot, Melanose, Pink Pitting, Scab	1.5	28.6	0.28	7	Begin applications when conditions favor disease development. Repeat sprays at 30 day intervals depending on disease severity.
Cucumber	Angular Leaf Spot, Downy Mildew	0.5 – 1.5 **	11.9	0.12	5	Apply weekly when plants begin to vine.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	0.5 – 1.5	17.9	0.18	7	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals depending on disease severity.
Pepper	Bacterial Spot	0.5 – 2	26.9	0.27	3	Begin applications when conditions first favor disease development and repeat at 3 to 10 day intervals depending on disease severity.
Tomato Processing	Anthracnose, Bacterial Speck, Bacterial Spot,	0.47 – 1.125	39.5	0.39	3	Begin applications when disease first threatens and repeat at 3 to 10 day

CROP	DISEASE	APP. RATE (TBSP PRODUCT / 1000 SQ. FT.)	MAX. APP. RATE/YEAR (TBSP Product /1000 SQ. FT.)	MAX. ANNUAL RATE (pounds Cu/1000 SQ. FT.)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Tomato Fresh market	Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	0.47 - 1.125	18.2	0.18		intervals depending on disease severity.
* Rosemary and Mint - In California, use the low rate of 0.75 lbs/A only.						
** Use the higher rates when conditions favor disease development.						

CONIFERS

For use on conifers, including Douglas Fir, Fir*, Juniper, Leyland Cypress*, Pine* and Spruce*, in Christmas tree plantings, forest stands and silviculture nurseries. For control of foliar diseases, apply BADGE X₂ as a thorough cover spray at rates ranging from 0.75 to 1.75 lbs/A. Begin applications in the spring at the initiation of new growth and repeat at 2 to 4 week intervals. Use the higher rates when disease pressure is severe or when environmental conditions favor disease development. **There is a maximum application rate of 2.0 lbs Cu/A with a maximum annual rate of 20 lbs Cu/A with a minimum retreatment interval of 7 days.**

BADGE X₂ may be used on the listed conifers for control of the following diseases:

CROP	LATIN NAME	DISEASE
Douglas Fir	<i>Pseudotsuga menziesii</i>	Rhabdocline Needlecast
Fir*	<i>Abies</i> spp.	Needlecasts
Juniper	<i>Juniperus</i> spp.	Anthrachnose, Phomopsis Twig Dieback*
Leyland Cypress*	<i>X Cupressocyparis leylandii</i>	Cercospora Needle Blight
Pine*	<i>Pinus</i> spp.	Needlecasts
Spruce*	<i>Picea</i> spp.	Needlecasts

Lichens*: To control lichens on any of the conifers above, apply 3.5 lbs of BADGE X₂ /A as a dormant application before new growth emerges in the spring. The addition of a non-ionic surfactant (NIS) will improve control. A second application may be required after 12 months.

NOTE: Do not buffer or combine with emulsifiable concentrate insecticides.

* Not for use in California

ORNAMENTALS

Use BADGE X₂ for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shadehouses, outdoor nurseries, and outdoor landscape plantings.

For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 1.5 to 5 lbs/A of BADGE X₂. When new growth is present, apply as a thorough cover spray at rates ranging from 1.5 to 2 lbs/A of BADGE X₂. **One and a half (1.5) level tablespoons of BADGE X₂ per 1000 square feet is equivalent to 2.4 lbs/A.** Begin application at first sign of disease and repeat at 7 to 14 day intervals; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist.

Unless otherwise noted, the maximum single application rate is 2 lbs of Cu/A and the maximum annual rate is 20 lbs of Cu/A. The minimum retreatment interval is 7 days.

BADGE X₂ may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to BADGE X₂ have been found to be acceptable for the specific genera and species listed on this label under the conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants and the wide range of growing conditions, it is impossible to test every one for sensitivity to BADGE X₂. Neither the manufacturer nor the seller has determined whether or not BADGE X₂ can be safely used on ornamental or nursery plants not listed on this label. The user should determine if BADGE X₂ can be used safely prior to commercial use. In a small area, apply the labeled rates to the plants in question, (bedding plants, foliage, etc.), and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use.

NOTE: This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, or other metallic surfaces.

ORNAMENTALS		
CROP	SCIENTIFIC NAME	DISEASE
Aglaonema*	<i>Aglaonema</i> spp.	Bacterial Leaf Spot
Althea (Rose of Sharon)	<i>Hibiscus syriacus</i>	Bacterial Leaf Spot
Andromeda, Japanese*	<i>Pieris japonica</i>	Leaf Spots, Twig Blight

ORNAMENTALS

CROP	SCIENTIFIC NAME	DISEASE
Aralia	<i>Dizygotheca elegantissima</i>	Alternaria, Cercospora Leaf Spot, Xanthomonas Leaf Spot
Arborvitae	<i>Thuja</i> spp.	Alternaria Twig Blight, Cercospora Leaf Blight
Aster*	<i>Aster</i> spp.	Downy Mildew, Leaf Spots
Azalea ¹	<i>Rhododendron</i> spp.	Botrytis Blight, Bud Blight*, Cercospora Leaf Spot, Phytophthora Dieback, Powdery Mildew, Twig Blight*
Beech*	<i>Fagus</i> spp.	Leaf Spots
Begonia	<i>Begonia semperflorens</i>	Bacterial Leaf Spot (<i>Erwinia</i> spp., <i>Pseudomonas</i> spp., <i>Xanthomonas</i> spp.)
Bougainvillea	<i>Bougainvillea spectabilis</i>	Anthrachnose, Bacterial Leaf Spot
Boxwood*	<i>Buxus</i> spp.	Leaf Spots
Camellia	<i>Camellia japonica</i> , <i>C. sasanqua</i>	Anthrachnose, Bacterial Leaf Spot
Camphor Tree	<i>Cinnamomum camphora</i>	Pseudomonas Leaf Spot
Canna	<i>Canna</i> spp.	Pseudomonas Leaf Spot
Carnation ¹	<i>Dianthus</i> spp.	Alternaria Blight, Botrytis Blight, Pseudomonas Leaf Spot
Cedar*	<i>Cedrus</i> spp.	Tip Blight
Cherry, Nanking*	<i>Prunus tomentosa</i>	Bacterial Leaf Spot
Chinese Tallow Tree	<i>Sapium sebiferum</i>	Bacterial Leaf Spot (<i>Pseudomonas</i> spp., <i>Xanthomonas</i> spp.)
Chrysanthemum ¹	<i>Dendranthema x grandiflorum</i>	Botrytis Blight, Pseudomonas Leaf Spot, Septoria Leaf Spot
Cotoneaster	<i>Cotoneaster</i> spp.	Botrytis Blight
Crabapple*	<i>Malus</i> spp.	Fire Blight
Cypress*	<i>Cupressaceae</i> spp.	Twig Blight
Dahlia	<i>Dahlia pinnata</i>	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Delphinium*	<i>Delphinium</i> spp.	Leaf Spots
Dianthus	<i>Dianthus</i> spp.	Bacterial Soft Rot, Bacterial Spot
Dogwood, Flowering	<i>Cornus florida</i>	Anthrachnose
Dogwood, Kousa*	<i>Cornus kousa</i>	Fungal Leaf Spot
Douglas Fir	<i>Pseudotsuga menziesii</i>	Rhabdocline Needlecast
Dracaena*	<i>Dracaena marginata</i>	Bacterial Leaf Spot
Dumb Cane*	<i>Dieffenbachia</i> spp.	Bacterial Leaf Spot
Dusty Miller	<i>Senecio cineraria</i>	Bacterial Leaf Spot (<i>Pseudomonas cichorii</i>)
Echinacea	<i>Echinacea</i> spp.	Bacterial Leaf Spot (<i>Pseudomonas cichorii</i>)
Elm, Chinese	<i>Ulmus parvifolia</i>	Xanthomonas Leaf Spot
Euonymus	<i>Euonymus</i> spp.	Anthrachnose, Botrytis Blight
Fern, Boston*	<i>Nephrolepis exaltata</i>	Bacterial Leaf Spot
Fern, Holly	<i>Crytomium falcatum</i>	Pseudomonas Leaf Spot
Fig, Weeping*	<i>Ficus benjamina</i>	Bacterial Leaf Spot
Filbert (Ornamental)*	<i>Corylus</i> spp.	Filbert Blight
Fir*	<i>Abies</i> spp.	Needlecasts
Gardenia	<i>Gardenia jasminoides</i>	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	<i>Pelargonium</i> spp.	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Gladiola	<i>Gladiolus</i> spp.	Alternaria Leaf Spot, Anthrachnose, Bacterial Leaf Blight, Botrytis; Gray Mold
Golden Rain Tree	<i>Koelreuteria paniculata</i>	Bacterial Leaf Spot
Grape Ivy*	<i>Cissus</i> spp.	Bacterial Leaf Spot
Hawthorn*	<i>Crataegus</i> spp.	Fire Blight
Hibiscus ⁴	<i>Hibiscus</i> spp.	Bacterial Leaf Spot
Holly*	<i>Ilex</i> spp.	Bacterial Blight, Leaf Spots
Honeylocust*	<i>Gleditsia triacanthos</i>	Bacterial Leaf Spot
Honeysuckle, Tatarian*	<i>Lonicera tatarica</i>	Bacterial Leaf Spot
Impatiens	<i>Impatiens wallerana</i>	Bacterial Leaf Spot
Indian Hawthorn ⁵	<i>Raphiolepis indica</i>	Anthrachnose, Entomosporium Leaf Spot
Iris ^{6*}	<i>Iris</i> spp.	Bacterial Leaf Spot
Ivy (English, Algerian) ¹	<i>Hedera helix</i> , <i>H. canariensis</i>	Xanthomonas Leaf Spot
Ixora	<i>Ixora coccinea</i>	Xanthomonas Leaf Spot
Juniper	<i>Juniperus</i> spp.	Anthrachnose, Phomopsis Twig Dieback*
Lantana	<i>Lantana camara</i>	Bacterial Leaf Spot
Leyland Cypress*	<i>X Cupressocyparis leylandii</i>	Cercospora Needle Blight
Lilac	<i>Syringa</i> spp.	Cercospora Leaf Spot, Pseudomonas Blight*
Lily, Easter ²	<i>Lilium longiflorum</i>	Botrytis Blight
Linden*	<i>Tilia</i> spp.	Anthrachnose, Leaf Blight
Loblolly Bay	<i>Gordonia lasianthus</i>	Anthrachnose
Loquat (non-bearing)	<i>Eriobotrya japonica</i>	<i>Colletotrichum</i> spp., <i>Entomosporium maculata</i>
Magnolia (Southern)	<i>Magnolia grandiflora</i>	Algal Leaf Spot, Anthrachnose, Bacterial Leaf Spot

ORNAMENTALS		
CROP	SCIENTIFIC NAME	DISEASE
Magnolia (Sweet Bay)	<i>Magnolia virginiana</i>	Anthraco
Magnolia (Oriental)	<i>Magnolia x soulangeana</i>	Bacterial Leaf Spot
Mandevilla	<i>Mandevilla</i> spp.	Anthraco
Maple*	<i>Acer</i> spp.	<i>Pseudomonas</i> Leaf Blight, Tar Leaf Spot
Marigold	<i>Tagetes</i> spp.	<i>Alternaria</i> Leaf Spot, <i>Botrytis</i> Leaf Rot, <i>Cercospora</i> Leaf Spot, Flower Rot
Mountain Ash*	<i>Sorbus</i> spp.	Fire Blight
Mulberry, Contorted*	<i>Morus bombycis</i>	Bacterial Leaf Spot
Mulberry, Weeping	<i>Morus alba</i>	Bacterial Leaf Spot
Narcissus*	<i>Narcissus</i> spp.	Leaf Blight
Nephtytis*	<i>Syngonium podophyllum</i>	Bacterial Leaf Spot
Oak*	<i>Quercus</i> spp.	Anthraco, Leaf Spots
Oak, Laurel	<i>Quercus laurifolia</i>	Algal Leaf Spot (<i>Cephaleuros virescens</i>)
Oleander	<i>Nerium oleander</i>	Bacterial Leaf Spot, Fungal Leaf Spot
Oregon Grapeholly*	<i>Mahonia aquifolium</i>	Leaf Spots
Pachysandra	<i>Pachysandra procumbens</i>	Canker*, Leaf Spots, Twig Blight*, <i>Volutella</i> Leaf Blight
Palm, Date (non-bearing)	<i>Phoenix dactylifera</i>	<i>Pestalotia</i> Leaf Spot
Palm, European Fan	<i>Chamaerops humilis</i>	<i>Pestalotia</i> Leaf Spot
Palm, Parlor*	<i>Chamaedorea elegans</i>	Bacterial Leaf Spot
Palm, Queen	<i>Arecastrum romanzoffianum</i>	<i>Exosporium</i> Leaf Spot, <i>Phytophthora</i> Bud Rot
Palm, Washingtonia	<i>Washingtonia robusta</i>	<i>Pestalotia</i> Leaf Spot
Peach (Flowering) (non-bearing) 3*	<i>Prunus</i> spp.	Bacterial Blast, Brown Rot, Fire Blight
Pear (Flowering) (non-bearing)	<i>Pyrus calleryana</i>	Fire Blight, Leaf Spot
Pentas (Egyptian Star)	<i>Pentas</i> spp.	Bacterial Leaf Spot (<i>Pseudomonas</i> spp.*, <i>Xanthomonas</i> spp.)
Peony	<i>Paeonia</i> spp.	<i>Botrytis</i> Blight
Periwinkle	<i>Catharanthus roseus</i> , <i>Vinca</i> spp.	<i>Phomopsis</i> Stem Blight
Philodendron	<i>Philodendron selloum</i>	Bacterial Leaf Spot
Phlox	<i>Phlox</i> spp.	<i>Alternaria</i> Leaf Spot
Photinia (Red Tip)	<i>Photinia x fraseri</i> , <i>P. glabra</i>	Anthraco, <i>Entomosporium</i> Leaf Spot
Pine*	<i>Pinus</i> spp.	Needlecasts
Pistachio	<i>Pistacia chinensis</i>	Anthraco
Plantain Lily 6	<i>Hosta</i> spp.	Bacterial Leaf Spot
Plum (Flowering) (non-bearing) 3*	<i>Prunus</i> spp.	Bacterial Blast, Bacterial Leaf Spot, Brown Rot, Fire Blight
Pothos*	<i>Scindapsus</i> spp.	Bacterial Leaf Spot
Powder Puff Plant	<i>Calliandra</i> spp.	Bacterial Leaf Spot
Pyracantha	<i>Pyracantha</i> spp.	Fire Blight, Scab
Rhododendron	<i>Rhododendron</i> spp.	<i>Alternaria</i> Flower Spot
Rose 1	<i>Rosa</i> spp.	Black Spot, Powdery Mildew
Snapdragon	<i>Antirrhinum majus</i>	Anthraco, Dieback, Downy Mildew
Spathe Flower*	<i>Spathiphyllum</i> spp.	Bacterial Leaf Spot
Spiraea*	<i>Spiraea</i> spp.	Fire Blight
Spruce*	<i>Picea</i> spp.	Needlecasts
Sycamore	<i>Platanus</i> spp.	Anthraco, Leaf Spots*
Tulip	<i>Tulipa</i> spp.	Anthraco, <i>Botrytis</i> Blight
Umbrella Tree*	<i>Schefflera</i> spp.	Bacterial Leaf Spot
Verbena	<i>Verbena</i> spp.	<i>Xanthomonas</i> Leaf Spot
Viburnum	<i>Viburnum odoratissimum</i> , <i>V. suspensum</i> , <i>V. plicatum</i>	Anthraco
Viola (Pansy, Violet)	<i>Viola</i> spp.	Downy Mildew
Willow	<i>Salix</i> spp.	Anthraco
Yew*	<i>Taxus</i> spp.	Needle Blight
Yucca (Adam's Needle)	<i>Yucca</i> spp.	<i>Cercospora</i> Leaf Spot, <i>Septoria</i> Leaf Spot
Zinnia*	<i>Zinnia</i> spp.	Leaf Spots

ORNAMENTALS

CROP	SCIENTIFIC NAME	DISEASE
* Not for use in California		
1	Can cause discoloration of foliage and/or blooms on some varieties. To prevent residues on commercial plants, do not spray immediately before selling season.	
2	Apply at 4.5 to 7.5 lbs/A. The maximum single application rate is 2.5 pounds of Cu per acre. The maximum amount of metallic copper which may be applied in a 12 month period is 75 pounds of Cu per acre. Do not apply any additional copper pesticide to this land for 36 months.	
3	Apply dormant through bloom only.	
4	Hibiscus - Do not apply to plants in flower.	
5	For Indian Hawthorne use 3 to 6 lbs/A.	
6	Some cultivars may be sensitive to BADGE X ₂ .	
NOTE: Phytotoxicity may depend on varietal differences. If unfamiliar with the use of BADGE X ₂ , apply the recommended rate to a few plants and observe after 7 to 10 days for symptoms of phytotoxicity.		

Control of Ball Moss*, Spanish Moss* and Lichens* on Ornamentals and Shade Trees: Apply BADGE X₂ in early spring when trees are dormant. Apply 9 to 12 pounds of BADGE X₂ in 100 gallons of water, using 1 ½ gallons of spray per foot of tree height. Be sure to thoroughly wet ball moss tufts, Spanish moss or lichens. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months. Do not exceed 7.09 lbs/A per 12 month period.

Live oak – Apply 2.5 to 12 lbs of Badge X₂. Apply in the spring when ball moss is actively growing in 100 gallons water. Use 1.5 gallons spray per foot of tree height. Ensure ball moss tufts are thoroughly wetted. The addition of nonionic surfactant will improve control. A follow up application may be needed 12 months later. Do not exceed 7.09 lbs/A per 12 month period.

NOTE: BADGE X₂ may be injurious to some ornamental plants growing beneath the trees. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Cold Storage Protection for Dormant Rootstock*: To protect bare-root nursery trees from Phytophthora Crown Rot and Botrytis, use 4 to 6 pounds of BADGE X₂ per 100 gallons of water. Apply as a dip or spray to the roots and lower stems of dormant rootstock prior to placing in cold storage. Do not apply to rootstock less than 2 years old.

*Not for use in California

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional Requirements for Aerial Applications

- The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter
- Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
- When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional Requirements for Ground Boom Application

- Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Chemigation Requirements

- Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Requirements for Chemigation Systems Connected to Public Water Systems

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favor drift beyond the area intended for treatment.

Requirements for Sprinkler Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed through storage and disposal.

Pesticide Storage

Store under well-vented, cool and dry storage conditions. Do not store under moist conditions.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling

Nonrefillable Paper and Plastic Bags

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration.

Nonrefillable Fiber Drums with Liners

Nonrefillable container. Do not reuse or refill this container. Completely empty the liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or return to the Seller, or dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Fiber drum is to be disposed of in the same manner required for its liner.

For rigid, non-refillable containers (2.5 to 5 gallons):

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of resulting smoke.

For rigid, non-refillable containers that are too large to shake (with capacities greater than 5 gallons):

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of resulting smoke.

Pressure Rinse Procedures (all sizes)

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire label before using this product, including this Limitation of Warranty and Liability.

If the terms are not acceptable, return the product at once unopened for a refund of the purchase price.

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use, subject to the inherent risks described below, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ISAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Buyers and Users of this product must be aware that there are inherent unintended risks associated to the use of this product, independent from the control of Isagro. These risks include, but are not limited to, weather conditions, soil factors, moisture conditions, diseases, irrigation practices, condition of the crop at the time of application, materials which are present in the tank mix with this product or prior to the application of it, cultural practices or the manner of use or application, all risks which are impossible to eliminate. The Buyers and Users should be aware that these factors may cause: ineffectiveness of the product, reduction of harvested yield of the crop (entirely or partially), crop injury or injury to non-target crops or plants or to rotational crops caused by carryover in the soil, resistance of the target diseases to this product. Therefore additional care, treatment and expense are required to take the crop to harvest.

If the Buyer does not agree with the acceptance of these risks, then THE PRODUCT SHOULD NOT BE APPLIED. To the extent consistent with applicable law, by applying this product the Buyer acknowledges and accepts these inherent unintended risks and AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

To the extent consistent with applicable law, in no event shall ISAGRO or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product (including claims based in contract, negligence, strict liability, other tort or otherwise). To the extent consistent with applicable law, the exclusive remedy of the User or Buyer and the exclusive Liability of Isagro or Seller shall be the return of the purchase price of the product, or at the election of Isagro or Seller, the replacement of the product.

To the extent consistent with applicable law, this Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

Isagro or its Seller must have prompt notice (within 7 days of observation) of any claim so that an immediate inspection of Buyer's or User's growing crops can be made. To the extent consistent with applicable law, if Buyer and User do not notify Isagro or Seller of any claims, in proper time, it shall be barred from obtaining any remedy.

To the extent consistent with applicable law, by applying this product the Buyers and Users are deemed to have accepted the terms of this Limitation of Warranty and Liability, which may not be modified by any verbal or written agreement.

Alliete and Roval are registered trademarks of Bayer CropScience.

Badge is a registered trademark of Isagro USA, Inc.

Curtec is a registered trademark of Curtec Corporation.

Kentan is a registered trademark of Isagro S.p.A. Corp.

Tre-Hold is a registered trademark of Amvac Chemical Corporation

AMEND 6April2015(e13April15)(26Aug2015)